MAPHEX MAnure Phosphorus EXtraction System



Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹



The Phosphorus Challenge

Phosphorus is a necessary and valuable nutrient

Manure has been valued for thousands of years

However:

To much manure P in areas of surfiet cause it to be a pollutant

P sorbing soil amendments and alternative application methods (injection, etc) are temporary solutions, at best.



But, what if manure P could be concentrated enough to be economically transported?

And, what if the liquid left behind had Negligible Phosphorus?

USDA

Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹



¹ USDA Agricultural Research Service, ² Pennsylvania State University - University Park, PA

Auger Press

Step 1: Initial Liquid/Solid Separation

Manure Slurry

Dairy manure slurry

Solids: 100%

Total P: 100%

Total P: 15%

Solids: Total p. 70.7%

Stackable
These low P solids a

Chemical Treatment

These low P solids are ideal for reuse as bedding material

Fe sulfate with Polymer

Chemical Treatment Tank

AutoVac®

Filtration

Total P: 45%
Stackable

Centrifuge

These high P solids are easily transported to areas with low P

Step 3:

Step 2:

Final Liquid/Solid Separation

This was manure slurry

10 minutes ago

Treated water is

Fine sized solids

Auger press


Total P: 40% Stackable

These high P solids are easily transported to areas with low P

96 – 99% P removal efficiency 99% solids removal efficiency All solids stackable (~70% moisture) Most nitrogen is retained (N:P ~ 50:1) Ideal for fertigation of crops pH unchanged by process **HIGHLIGHTS**

Cost for a 1000 cow dairy
Per day \$750
per cow/yr \$180
per kg P removed \$38
per lb P removed \$17

Clinton D. Church, USDA-ARS, 3702 Curtin Road, University Park, PA 16802, cell 858.212.5104, Clinton.Church@ars.usda.gov






Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹





to areas with low P






Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹

¹ USDA Agricultural Research Service, ² Pennsylvania State University - University Park, PA



Step 2: Chemical Treatment





Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹

¹ USDA Agricultural Research Service, ² Pennsylvania State University - University Park, PA







Clinton D. Church¹, Alex N. Hristov², Ray B. Bryant¹, and Peter J. A. Kleinman¹

 1 USDA Agricultural Research Service, 2 Pennsylvania State University - University Park, PA





96 – 99% P removal efficiency 99% solids removal efficiency All solids stackable (~70% moisture) Most nitrogen is retained (N:P ~ 50:1) Ideal for fertigation of crops pH unchanged by process **HIGHLIGHTS**

Cost for a 1000 cow dairy	
Per day	\$750
_per cow/yr	\$180
per kg P removed	\$38
per lb P removed	\$17

Full-Scale MAPHEX System



Uses for Solids

- 1)Low P composted bedding from bulk solids
- 2) Sale of high P nutrient solids to organic farmers, mushroom growers, or retail
- 3) Feedstocks for energy generation
- 4) Innovative products (e.g. cowpots®)

Current Work

- 1) Testing of the Full-Scale Mobile Treatment System
- 2) Ongoing work to lower daily operating costs